

MALAYSIAN JOURNAL OF ANALYTICAL SCIENCES

Published by The Malaysian Analytical Sciences Society

ISSN 1394 - 2506

THE CHEMICAL CONSTITUENTS FROM THE FRUITS OF Johannesteijsmannia altifrons AND ANTIBACTERIAL ACTIVITY

(Komposisi Kimia Daripada Buah Johannesteijsmannia altifrons dan Aktiviti Antibakteria)

Noor Aziiraa Sabri¹, Wan Yaacob Wan Ahmad¹*, Nur Shafiqa Abdullah¹, Nazlina Ibrahim²

¹School of Chemical Sciences and Food Technology, Faculty of Science and Technology ²School of Biosciences and Biotechnology, Faculty of Science and Technology Universiti Kebangsaan Malaysia, 43600 UKM Bangi, Selangor, Malaysia

*Corresponding author: wanyaa@ukm.edu.my

Received: 10 October 2016; Accepted: 16 January 2017

Abstract

Isolation on the hexane extract of the fruits of *Johannesteijsmannia altifrons* by vacuum liquid and radial chromatography yielded three known compounds namely β -sitosterol, γ -taraxasterol and stigmasterol. Their structures were determined using one-dimensional and two-dimensional nuclear magnetic resonance spectroscopy, fourier transform infra-red spectroscopy and mass spectrometry. All the isolated compounds were tested for their antibacterial activity using disc diffusion and minimum inhibitory concentration (MIC). All three compounds showed weak and moderate activities against all nine tested bacteria. This is the first report on the isolation of compounds from this species and on the antibacterial activity of γ -taraxasterol.

 $\textbf{Keywords}: \textit{ Johannesteijs mannia altifrons}, \beta \text{-sitosterol}, \gamma \text{-taraxasterol}, \text{ stigmasterol}, \text{ antibacterial}$

Abstrak

Pemencilan ke atas ekstrak heksana bagi buah *Johannesteijsmannia altifrons* dengan kromatografi cecair vakum dan radial menghasilkan tiga sebatian yang pernah ditemui bernama β -sitosterol, γ -taraksasterol dan stigmasterol. Strukturnya ditentukan dengan spektroskopi resonans magnet nukleus satu-dimensi dan dua-dimensi, spektroskopi infra-merah transformasi Fourier dan spektrosmetri jisim. Kesemua sebatian yang dipencilkan diuji aktiviti bakterianya dengan kaedah peresapan cakera dan kepekatan perencatan minimum (KPM). Kesemua tiga sebatian menunjukkan aktiviti yang lemah dan sederhana terhadap kesemua sembilan bakteria yang diuji. Ini merupakan laporan pertama mengenai pemencilan sebatian daripada spesies ini dan mengenai aktiviti antibakteria bagi γ -taraksasterol.

Kata kunci: *Johannesteijsmannia altifrons*, β-sitosterol, γ-taraksasterol, stigmasterol, antibakteria

Introduction

Johannesteijsmannia belonging to the Arecaceae family contains four species which are J. altifrons, J. lanceolata, J. magnifica and J. perakensis. Johannesteijsmannia altifrons which is also known as "Daun Sang" (Malay) and Joey Palm (English) commonly found in Kelantan, Perak, Terengganu, Pahang, Johor and Kedah [1]. No chemical constituents have ever been isolated or characterized from Johannesteijsmannia altifrons. The family of Arecaceae has been reported to contain flavonoids [2], alkaloids [3], terpenoids, steroids, fatty acids and tannins [4]. Thus, conducting a phytochemical study on Johannesteijsmannia altifrons is vital to further enrich our knowledge on the species. In this study, the hexane extract from fruits of Johannesteijsmannia altifrons was purified and led to the discovery of known β-sitosterol, γ-taraxasterol and stigmasterol. Previously, Singh and Ghanapriya [5] reported β-sitosterol and stigmasterol inhibited at low levels (1+) against Staphylococcus aureus, Bacillus subtilis and

Escherichia coli. According to previous study by Manriquez et al. [6], both β - sitosterol and stigmasterol gives MIC value at 16.5 μg/mL against Staphylococcus aureus. Previous study discovered no inhibition of β -sitosterol and stigmasterol reported on Escherichia coli, Bacillus subtilis and Salmonella thyphimurium [6]. There are three types of extracts in this study which are hexane, chloroform and methanol but hexane extract was chosen to be purified further. This is because compounds from hexane extract are easily separated and purified compared to the other two extracts.

Materials and Methods

General

Different spectroscopic methods were used to elucidate the structures of β -sitosterol, γ -taraxasterol and stigmasterol including ultraviolet–visible spectroscopy, mass spectroscopy, fourier transform infra-red spectroscopy, one-dimensional, two-dimensional and carbon-13 nuclear magnetic resonance spectroscopy. The one-dimensional, two-dimensional and carbon-13 nuclear magnetic resonance spectroscopy spectra were recorded using Avance III 600 MHz Bruker. Deuterated chloroform (CDCl₃) was used as the NMR solvent with the chemical shifts, δ_H , in ppm and the values of coupling constants, J, in Hz. Mass spectra were recorded by using the LC-mass spectrometry (LC-MS) on the Dionex/Bruker Micro ToFQ. The infra-red spectra were recorded on the Perkin-Elmer Spectrum 400 and the UV-VIS spectra were obtained in chloroform using UV-2400PC. Vacuum liquid chromatography (VLC) was prepared by using silica gel of Merck 7747. For radial chromatography (RC), silica gel with PF₂₅₄ containing gypsum (Merck 7749) was used. Thin layer chromatography (TLC) was performed on commercially pre-coated aluminum silica gel 60 F₂₅₄ of Merck 5554.

Plant materials

The fruits of *Johannesteijsmannia altifrons* were collected at Pelagat Forest Reserve, Besut, Terengganu in February 2015. A voucher specimen (UKMB 40309) was deposited at the Herbarium of Universiti Kebangsaan Malaysia, Bangi (UKMB).

Extraction and isolation

The air-dried, fruits of *Johannesteijsmannia altifrons* (786 g) were extracted for 3 days in methanol at room temperature, followed by filtration to give methanol solution. They were then extracted with n-hexane three times, combined and rotary evaporated to afford 4.42 g (0.56%) of greenish brown fruit n-hexane extract. The extract was fractionated by VLC, eluted with increasing polarity of n-hexane-ethyl acetate to collect the less polar and more polar fractions. The fractions were combined based on their silica gel TLC profiles. The VLC gave 16 fractions (F₁- F₁₆). Since fractions F₄ to F₇ have similar components, they were combined and further fractionated using RC to produce four combined fractions (F₄₁-F₄₄). The third fraction (F₄₃) contained β -sitosterol (32 mg). Then, fractions F₉-F₁₂ were combined and purified by RC to yield 7 combined fractions (F₉₁-F₉₇); fractions F₉₄ and F₉₆ contained γ -taraxasterol (25 mg) and stigmasterol (0.55 g), respectively.

Bioassay

The pure compounds were assayed for antibacterial activity by using two methods which are disc diffusion and minimum inhibitory concentration (MIC) as described by Mamman et al. [7], Mukhtar and Ghori [8], and Yap et al. [9] with some modification, against nine bacteria; *Staphylococcus aureus*, Methicillin-resistant *Staphylococcus aureus* (MRSA), Bacillus subtilis, Bacillus thruriengiensis, Escherichia coli, Salmonella thyphimurium, Serratia marcescens, Vibrio fluvialis and Vibrio cholera. The pure compounds were prepared at 30 µg/mL in DMSO.

Disc diffusion

Disc diffusion method for antibacterial testing was carried out to determine antibacterial activity of the pure compounds. The nine bacteria were spread on the Mueller-Hinton agar surface by using a sterile swab and the plates were dried for 15 minutes. Then, $100~\mu L$ of each dilution of the pure compounds was impregnated into sterile, blank disc of 6 mm in diameter and placed on the Mueller-Hinton agar surface. The positive control used was chloramphenicol 30 μg /disc which is a standard commercial antibiotic. The negative control used was DMSO (100%). The plates were then incubated at 37 °C for 24 hours. After the incubation, the plates were examined for inhibition zone and the results were recorded. The diameter of inhibition zones were measured in millimeter (mm). The test was repeated three times to ensure reliability.

Minimum inhibition concentration

Minimum Inhibition Concentration (MIC) was carried out to determine the lowest concentration of the compounds that could inhibit the visible growth of tested bacteria by using 96 microwell plate. An amount 100 μ L of pure compounds undergo dilutions from 30 to 0.117 μ g/mL. 50 μ L of nine bacteria were added into wells containing compounds. The 96 microwell plate was incubated at 37 °C for 24 hours. The positive and negative controls used were 100 μ L of respective chloramphenical antibiotic and DMSO. Then, 50 μ L of 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) was added to each single well and the plate was incubated at 37 °C for 2 hour. The colour development was observed and the results were recorded. The test was repeated three times to ensure reliability.

Results and Discussion

The ESI-MS gave molecular ion at $[M+Na]^+$ m/z 437.3381 which β-Sitosterol appeared as colourless needles. The ESI-MS gave molecular ion at $[M+Na]^+$ m/z 437.3381 which corresponded to the molecular formula $C_{29}H_{50}O$ (414.3862 + 22.9898 = 437.3759; Δ 0.40513). The IR spectrum showed absorption peaks at 3385 cm⁻¹ which is characteristic of O-H stretching, peaks at 2936 and 2868 cm⁻¹ due to aliphatic or C-H stretching (CH₃). Other absorption frequencies include peak at 1708 cm⁻¹ due to absorption of C=C stretching. The 1H NMR spectrum of β-sitosterol showed a multiplet signal at δ_H 3.53 (H-3), indicates the presence of a hydroxyl methine proton. The occurrence of a highly-deshielded signal at δ_H 5.36 (H-6) indicates the presence of a vinylic proton. On the other hand, a singlet signal at δ_H 1.02 indicates a methyl group at H-29. Two doublet signals at δ_H 0.86 and 0.91 are due to the two non-equivalent methyls at H-26 and H-27 of an isopropyl group. The ^{13}C NMR spectrum showed that β -sitosterol contains 29 different carbon atoms consisting of six methyls, 11 methylenes, nine methines, and three quaternary carbons. Signal of methine carbon at δ_C 71.8 ppm indicates a carbon attached to an oxygen atom which showed the existence of a hydroxyl group at C-3. Two highly deshielded signals at δ_C 140.8 and 121.7 ppm, which are assigned to C-5 and C-6 indicates there is one carbon double bond in the molecule. According to the available data as well as comparison with the literature [10], the compound is confirmed to be β -sitosterol.

γ-Taraxasterol was obtained as colourless needle crystal. The ESI-MS showed a molecular ion peak at [M+Na]⁺ m/z 449.2842 which corresponded to the molecular formula $C_{30}H_{50}O$ (426.3862 + 22.9898 = 449.3759; Δ 0.09176). The IR spectrum displayed a broad absorbance peak at 3385 cm⁻¹ which characteristic of O-H stretching. Absorptions at 2927 and 2871 cm⁻¹ represent the Csp^3 -H stretching. The ¹H NMR spectrum showed signal of a proton at δ_H 3.51 (H-3) is attached to the oxygen-binding carbon (C-3). This is deshielded proton which is hydroxyl methine proton. Seven singlet methyls at signals δ_H 1.61 (H-23), 1.10 (H-24), 0.82 (H-25), 0.81 (H-26), 1.29 (H-27), 0.80 (H-28) and 1.05 (H-30) indicating that they are attached to quaternary carbons and bound directly to the ring system. There is only one highly deshielded proton at δ_H 5.37 (H-21) indicating the olefinic proton that attaches to a double bond carbon (C-21). The ¹³C NMR spectrum showed γ-taraxasterol is formed by 30 carbon atoms consisting of eight methyls, nine methylenes, seven methines, and six quarternary carbons. A methine carbon at δ_C 80.7 indicates a carbon attached to an oxygen atom which showed the existence of a hydroxyl group at C-3. There are two highly deshielded olefinic carbon signals at δ_C 141.0 (C-20) and 121.7 (C-21) suggests that there is one double bond contained in the molecule. Through comparison of the observed data with literature [11], the compound is identified as γ-taraxasterol.

Stigmasterol was isolated as colourless needles. The ESI-MS gave molecular ion at $[M+Na]^+$ m/z 435.3279 which corresponded to the molecular formula $C_{29}H_{48}O$ (412.3705 + 22.9898 = 435.3603; Δ 0.0324). The IR spectrum displayed stretching band for O-H at 3354 cm⁻¹. Peaks at 2934 and 2866 cm⁻¹ indicate the presence of aliphatic or C-H stretching (CH₃). A band at 1667 cm⁻¹ appears due to the existence of olefinic group in this compound. The ¹H NMR spectrum displayed signals for six methyls. Two singlet methyls at signals δ_H 0.69 (H-18) and 1.02 (H-19) indicating that they are bound directly to the ring system. Three doublet methyls at δ_H 0.93 (H-21), 0.84 (H-26) and 0.7 (H-27), and one *triplet* methyl at δ_H 0.83 (H-29) are located at the side chain outside the rings of the main skeleton. Hydroxyl methine proton appeared at δ_H 3.54 (H-3) which attached to the oxygen-binding carbon (C-3). Three highly-deshielded protons presence at δ_H 5.36, 5.03, and 5.09 due to the presence of two pairs of double bond which assigned at H-6, H-22 and H-23. The ¹³C NMR spectrum showed stigmasterol contained 29 carbon atoms consisting of six methyls, nine methylenes, 11 methines, and three quaternary carbons. Signal at δ_C 71.8 indicates a

methine carbon attached to an oxygen atom which showed the existence of a hydroxyl group at C-3. There are four olefinic carbon signals that appear downfield and highly deshielded at δ_C 140.8 (C-5), 121.7 (C-6), 138.3 (C-22) and 129.3 (C-23) suggests that there are two double bonds in the molecule. Based on the discussion above and data comparison with literature [10], the compound is identified as stigmasterol. The spectra of β -sitosterol, γ -taraxasterol and stigmasterol are shown in Figures 1, 2 and 3, respectively.




Figure 1. The infra-red spectrum of β -sitosterol




Figure 2. The infra-red spectrum of γ -taraxasterol




Figure 3. The infra-red spectrum of stigmasterol

β-Sitosterol
$$\gamma$$
-Taraxasterol γ -Taraxasterol γ -Taraxasterol γ -Taraxasterol

Figure 4. Structures of the pure compounds

Antibacterial activity

The pure compounds isolated from the fruits of *Johannesteijsmannia altifrons* were tested for antibacterial activity. In disc diffusion method, β -sitosterol showed low inhibition which is 7.0 ± 0.00 mm against *Vibrio fluvialis*, MRSA and *Bacillus subtilis* whereas γ -taraxasterol and stigmasterol exhibited moderate activities against *Escherichia coli*, *Serratia marcescens*, *MRSA* and *Bacillus subtilis*. Based on the results, it was observed that β -sitosterol has inhibition against one gram negative bacteria and two gram positive bacteria. This may probably due to the morphological differences between gram negative and gram positive bacteria in terms of their cell wall composition. The gram positive bacteria only have an outer peptidoglycan which is not effective permeability barrier compared to gram negative bacteria that have both outer membrane and peptidoglycan. This makes the gram negative bacteria cell wall impermeable to most of the drugs [12]. For stigmasterol, the inhibition occurs at low level (1+) against *Staphylococcus aureus*, *Bacillus subtilis* and *Escherichia coli* which is similar as reported by Singh dan Ghanapriya [5]. Besides, γ -taraxasterol showed moderate (2+) inhibition against *Serratia marcescens* and *Bacillus subtilis*. Kanamysin and vancomysin used as positive controls and the results showed all compounds have lower inhibitory activity against all tested bacteria compared to the antibiotics. DMSO as negative control did not show any antibacterial activity against tested bacteria. The inhibitory zone values for all compounds are shown in Table 1.

Table 1. Antibacterial activity of the pure compounds in disc diffusion method

Pure Compounds	β-Sitosterol	γ-Taraxasterol	Stigmasterol	Kanamysin (K)*/ Vancomysin (V)**	
	Inhibitory Zone (mm)				
Escherichia coli *	-	7.0 ± 0.53	7.0 ± 0.00	$20 \pm 1.00 (K)$	
Vibrio fluvialis *	7.0 ± 0.00	7.0 ± 0.00	7.0 ± 0.00	35 ± 1.04 (K)	
Salmonella thyphimurium *	-	7.0 ± 0.00	7.0 ± 0.00	$26 \pm 0.00 (K)$	
Vibrio cholera *	-	-	-	$10 \pm 0.00 (K)$	
Serratia marcescens *	-	11.0 ± 0.00	11.0 ± 0.00	$23 \pm 1.52 (K)$	
Methicillin-resistant Staphylococcus aureus (MRSA) ***	7.0 ± 0.00	7.0 ± 0.00	8.0 ± 0.5	$20 \pm 0.00 \text{ (V)}$	
Bacillus thruriengiensis **	_	-	-	$22 \pm 0.00 (V)$	
Bacillus subtilis **	7.0 ± 0.00	10.0 ± 0.5	7.0 ± 0.00	$28 \pm 0.00 (V)$	
Staphylococcus aureus **	-	7.0 ± 0.53	7.0 ± 0.00	$25 \pm 0.00 \text{ (V)}$	

Mean measurement: SD (n = 3), *Gram negative, ** Gram positive

In minimum inhibition concentration (MIC) method, all compounds gave value of 7.5 μ g/mL against Staphylococcus aureus, Escherichia coli, Vibrio cholera, Serratia marcescens, Bacillus thruriengiensis and Bacillus subtilis whereas previous report showed both β -sitosterol and stigmasterol have higher MIC value of 16.5 μ g/mL and no inhibition against Escherichia coli and Bacillus subtilis. No explanation can be given on the difference. First report on antibacterial activity of γ -taraxasterol provides new knowledge for the compound. The MIC values are shown in Table 2.

Table 2. Antibacterial activity of the pure compounds in minimum inhibitory concentration (MIC) method

Pure compounds	β-Sitosterol	γ-Taraxasterol MIC (μg/mL)	Stigmasterol
Escherichia coli	7.5	7.5	7.5
Vibrio fluvialis	30	30	30
Salmonella thyphimurium	15	15	15
Vibrio cholera	7.5	7.5	7.5
Serratia marcescens	7.5	7.5	7.5
Methicillin-resistant Staphylococcus aureus (MRSA)	15	15	30
Bacillus thruriengiensis	7.5	7.5	7.5
Bacillus subtilis	7.5	7.5	7.5
Staphylococcus aureus	7.5	7.5	7.5

Conclusion

The investigation on the chemical constituents from fruits of *Johannesteijsmannia altifrons* resulted in the isolation of three known compounds which are β -sitosterol, γ -taraxasterol and stigmasterol. In disc diffusion method, the isolated compounds showed no inhibition against *Vibrio cholera* and *Bacillus thruriengiensis*, and weak or moderate activities against other seven bacteria. In MIC method, the lowest and highest MIC values were at respective 7.5 and 30 μ g/mL. All the compounds have been reported for the first time from this plant.

Acknowledgment

Authors would like to thank the Ministry of Education, Malaysia and Universiti Kebangsaan Malaysia (UKM) for the financial support under DLP-2013-018 grant and "Program Pelajar Cemerlang (PPC)" scholarship under Public Service Department, Malaysia. Special thanks to School of Chemical Sciences and Food Technology, Faculty of Science and Technology, UKM and Centre for Research and Innovation Management (CRIM) UKM for technical support on NMR, LC-MSToF, UV and FTIR instrumentation.

References

- 1. Dransfield, J. (1972). The genus *Johannesteijsmannia* H. E. Moore Jr. *The Gardens Bulletin Singapore*, 26: 63–83.
- 2. Muhaisen, H. (2002). Flavonoids from the base leaves of *Caryota urens* (Palmae). *American Scientific Publishers*, 6(11): 1225 1229.
- 3. Dyana, J. P. and Kanchana, G. (2012). Preliminary phytochemical screening of *Cocos nucifera* L. flowers. *International Journal of Current Pharmaceutical Research*, 4(3): 3 4.
- 4. Benmehdi, H., Hasnaoui, O., Benali, O. and Salhi, F. (2012). Phytochemical investigation of leaves and fruits extracts of *Chamaerops humilis* L. *Journal of Material and Environmental Science*, 3(2): 320 237.
- 5. Singh, S. and Ghanapriya, N. (2014). Chemical constituents and antibacterial activity of *Phaleria macrocarpa*. *International Journal of Pharmaceutical Sciences and Research*, 5(8): 3157 3162.
- 6. Manríquez-Torres, J. J., Zúñiga-estrada, A., González-ledesma, M. and Torres-valencia, J. M. (2007). The antibacterial metabolites and proacacipetalin from *Acacia cochliacantha*. *Journal of the Mexican Chemical Society*, 51(4): 228 231.
- 7. Mamman, P. H., Mshelia, W. P., Susbatrus, S. C. and Sambo, K. W. (1989). Antibacterial effects of crude extract of *Azadirachta indica* against *Escherichia coli, Salmonella* 142 spp and *Staphylococcus aureus*. *International Journal of Medicine and Medical Sciences*, 5(1): 14 18.
- 8. Mukhtar, S. and Ghori, I. (2012). Antibacterial activity of aqueous and ethanolic extracts of garlic, cinnamon and turmeric against *Escherichia coli* ATCC 25922 and *Bacillus subtilis* DSM 3256. *International Journal of Applied Biology and Pharmaceutical Technology*, 3(2): 131 136.

- 9. Yap, P. S. X., Chong Y. T. E., Wong, Y. Y., Lim, X. Y., Ang J. H., Goh, H. S. L., Lai, M. X. S, Mai, C. W., Buru, S. A., Pichika, M. R. and Lim, S. H. E. (2013). Antibacterial and antifungal testing of the different extracts of *Dillenia obovata* (blume) Hoogl. *World Journal of Pharmacy and Pharmaceutical Sciences*, 2(5): 3946 3962.
- 10. Chaturvedula, V. S. P. and Prakash, I. (2012). Isolation of Stigmasterol and β-Sitosterol from the dichloromethane extract of *Rubus suavissimus*. *International Current Pharmaceutical Journal*, 1(9): 239 242.
- 11. Yekta, M. M., Hamid, S. and Alavi, R. (2008). New triterpenoids from *Peucedanum ruthenicum*. *Iranian Journal of Pharmaceutical Sciences*, 4(4): 289 294.
- 12. Ghansar, N. H., Bhopale, S. U. and Prabhune, V. S. (2012). *In vitro* studies of antimicrobial properties of extracts from Unani medicinal plants. *International Journal of Pharmaceutical Bio Sciences*, 3(1): 240 249.